

Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Br

Search PubMed for Limits Preview/Index History Clipboard Details

About Entrez

Display Abstract Show: 20 Sort Send to Text

Text Version

Entrez PubMed  
Overview  
Help | FAQ  
Tutorial  
New/Noteworthy  
E-Utilities


PubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources  
Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

Privacy Policy

1: Int J Radiat Oncol Biol Phys. 1998 Jan 15;40(2):331-6.

Related Article



## Integration of filgrastim into chemoradiation for limited small cell lung cancer: a Phase I study.

Glisson B, Komaki R, Lee JS, Shin DM, Fossella F, Murphy WK, Kurie Perez-Soler R, Schea R, Vadhan-Raj S.

Department of Thoracic/Head and Neck Medical Oncology, The University of Texas M. D. Anderson Cancer Center, Houston 77030, USA.

**PURPOSE:** Recent studies document the value of early combined modality of small cell lung cancer, but also indicate that early thoracic radiation adds to myelosuppression and can complicate further chemotherapy. Other studies indicate that simultaneous use of growth factors with thoracic radiation may be deleterious. However, temporal separation of growth factor use from cytotoxic therapy may allow dose intensity to be maintained/enhanced during combined modality treatment. We sought to integrate filgrastim into a novel chemoradiation regimen for patients with limited small cell lung cancer using an approach that separates growth factor administration from both chemotherapy and thoracic radiation. **METHODS AND MATERIALS:** Twenty-seven patients with limited disease small cell lung cancer were enrolled in a Phase I trial of cisplatin, ifosfamide/mesna, oral etoposide, and thoracic radiation (1.5 Gy b.i.d. x 30 fractions days 1-19 cycle 1) +/- filgrastim (5 microg/kg/day). Filgrastim was on days 20-25 of cycle 1 after completion of radiation and following completion of oral etoposide in subsequent cycles. The primary end point was determination of maximum tolerated dose (MTD) of chemotherapy. Serial cohorts were treated with and without filgrastim. **RESULTS:** Because of dose-limiting thrombocytopenia primarily, and nonhematologic toxicity, the MTDs with and without filgrastim were identical (cisplatin 20 mg/m<sup>2</sup> i.v. and ifosfamide 1200 mg/m<sup>2</sup> i.v., both days 1-3, and etoposide 40 mg/m<sup>2</sup> p.o. days 1-14). Filgrastim use shortened the duration of neutropenia at the MTD (median 4 vs. 7 days), but was not associated with a reduction in febrile neutropenia. Although growth factor administration did not allow dose escalation of this regimen, it did allow chemotherapy doses to be maintained at the MTD more frequently through four cycles of therapy. In the evaluable patients, the overall response rate was 100% (71% partial and 29% complete). **CONCLUSIONS:** Despite careful attention to the timing of growth factor with chemoradiation, the administration of filgrastim with this regimen did not allow dose escalation. As in many other recent studies of hematopoietic

factors given prophylactically with chemotherapy, the duration of neutropeni  
the MTD was shortened and the need for dose reduction throughout treatmen  
reduced in patients receiving filgrastim at the MTD.

Publication Types:

- Clinical Trial
- Clinical Trial, Phase I

PMID: 9457817 [PubMed - indexed for MEDLINE]

---

Display: Abstract Show: 20 Sort Send to: Text

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

Mar 11 2004